SYMPOSIUM

# Understanding Biological Roles of Venoms Among the Caenophidia: The Importance of Rear-Fanged Snakes

Stephen P. Mackessy[1] and Anthony J. Saviola

School of Biological Sciences, University of Northern Colorado, 501 20th St, Greeley, CO 80639-0017, USA

From the symposium ''Integrative and Comparative Biology of Venom'' presented at the annual meeting of the Society for Integrative and Comparative Biology, January 3–7, 2016 at Portland, Oregon.

[1]E-mail: stephen.mackessy@unco.edu

**Synopsis** Snake venoms represent an adaptive trophic response to the challenges confronting a limbless predator for overcoming combative prey, and this chemical means of subduing prey shows several dominant phenotypes. Many front-fanged snakes, particularly vipers, feed on various vertebrate and invertebrate prey species, and some of their venom components (e.g., metalloproteinases, cobratoxin) appear to have been selected for ''broad-brush'' incapacitation of different prey taxa. Using proteomic and genomic techniques, the compositional diversity of front-fanged snakes is becoming well characterized; however, this is not the case for most rear-fanged colubroid snakes. Because these species consume a high diversity of prey, and because venoms are primarily a trophic adaptation, important clues for understanding specific selective pressures favoring venom component composition will be found among rear-fanged snake venoms. Rear-fanged snakes typically (but not always) produce venoms with lower complexity than front-fanged snakes, and there are even fewer dominant (and, arguably, biologically most relevant) venom protein families. We have dem-

them to feed infrequently. Many snakes commonly consume prey weighing 20% of their own body mass, and some venomous snakes have been reported to subdue and consume prey that exceed their own body mass by 50% or more (Greene 1984, 1992). However, swallowing large, potentially dangerous prey necessitates subduing the quarry without sustaining injury in the process, and snakes have evolved various means to facilitate successful prey capture.

The vast majority of non-venomous snakes utilize constriction or body pinning to restrain prey, whereas other species simply employ a jaw-holding technique, or swallow prey whole with little to no body involvement (Greene 1997; Cundall and Greene 2000; Bealor and Saviola 2007). Coachwhips (*Ma. c h f.age* ), for example, subdue and consume rodent prey by pinning and swallowing them (Werler and Dixon 2000; Bealor and Saviola 2007); however, this behavior results in significantly longer capture latencies and higher frequencies of prey escape compared to constriction by non-venomous snakes (Bealor and Saviola 2007). In laboratory trials, multiple predatory strikes may be necessary for a non-venomous, non-constricting snake to grasp and subdue prey successfully, but this opportunity is highly unlikely in the wild. Constricting prey is undoubtedly advantageous, as multiple body loops provide more contact with prey and limit the opportunity of prey escape, but it still requires the snake to be in constant contact with prey and increases the likelihood of sustaining injury from the retaliating animal.

Venomous snakes, on the other hand, subdue prey with venom. This complex mixture of proteins and peptides has allowed for the trophic transition from a mechanical (body pinning and constriction) to a chemical (venom) means of immobilizing prey. Venoms exhibit tremendous diversity and may vary

significant quantities of venom rapidly. Instead, venom is introduced into tissue more slowly, although often at multiple sites, and rear-fanged snakes produce multiple puncture wounds by chewing on prey (see below for feeding strategies). A more recent study, however, has suggested that the grooved enlarged rear maxillary teeth (as seen in many *Boiga* species; Mackessy 2010) are capable of delivering venom effectively and rapidly into tissue (Young et al., 2011). Combined, these features result in a low-pressure venom delivery system that delivers significantly lower quantities of venom when compared to the majority of front-fanged species.

### Venom extraction

The clear distinction between high- and low-pressure venom delivery systems also requires two very different venom extraction methods. Front-fanged venomous snakes, with the compressor glandulae muscle surrounding the venom gland, and a large quantity of venom stored in the basal lumen, can be easily extracted by manual expression of the venom gland. We have found that anesthetizing front-fanged snakes with isoflurane prior to extraction provides an effective way for quickly manipulating snakes and extracting significant quantities of venoms, and this also minimizes the risk to the handler and the snake. However, venom extraction from rear-fanged venomous snakes is more challenging, time-consuming, and generally results in significantly lower venom yields. We routinely anesthetize rear-fanged snakes with ketamine (Hill and Mackessy 1997; Mackessy et al. 2006), although anesthetics such as Zoletile, Tiletamine, and Zolazepam have also been used (Fry et al., 2003a,b). To stimulate salivation once the animal is anesthetized, an injection of pilocarpine is administered, and venoms are obtained by placing glass micropipettes over the animal's posterior maxillary teeth. Extractions may take up to 30 to 60 min, and venom yields are often a fraction of what is obtained from a front-fanged snake. However, these methods, and especially the use of pilocarpine, have been shown to increase venom yields (Hill and Mackessy 1997; Mackessy et al. 2006; Ching et al. 2012), and provide a safe and effective method for venom extraction for both the handler and the snake.

## Venom composition

Snake venoms are complex mixtures of proteins and peptides that exhibit a myriad of biological effects (Mackessy 2010). The majority of venomous snakes are found in the families Colubridae, Elapidae, and Viperidae, and venoms from each of these families exhibit dramatically different compositions, while most species within each family share many toxin families. Although a given venom may contain up to 100 different proteins (including isoforms), the vast majority of venom compounds can be classified into approximately 24 distinct protein families (Table 1). Viperid venoms are rich in enzymes such as phospholipase $A_2$ ($PLA_2$

**Fig. 1** Simplified comparison of venom composition for colubroid snakes, emphasizing dominant protein families. (**A**) Several examples from major extant clades of snakes. Note that for each species, only a few protein families tend to predominate. (**B**) Variation in venom proteomes between major clades and within the rattlesnakes (*Crotalus* and *Sistrurus*). In the elapid and colubrid examples given, 3FTxs dominate the venom proteome. This toxin family is typically not expressed in viperid venoms. Rattlesnake venoms show two major patterns: type I, dominated by SVMPs and having lower toxicity, and type II, dominated by presynaptic neurotoxins (crotoxin, Mojave toxin, etc.) and having high toxicity. Diagrams derived from sources indicated. CTL, C-type lectin; MyoTx, myotoxin a; PLA

2016). SVMP, SP, and $PLA_2$ activities have been documented in many rear-fanged snake venoms (Mackessy 2002; Saviola et al. 2014), and several of these enzymes, as well as non-enzymatic CRISPs and 3FTxs, have been isolated and characterized from rear-fanged venoms (Fry et al. 2003a; Pawlak et al. 2006; Peichoto et al. 2007, 2009; Weldon and Mackessy 2012). However, rear-fanged venomous snakes represent very different evolutionary lineages from viperids and elapids (Pyron et al. 2013; Vidal 2002), and they comprise several families, subfamilies, and hundreds of species and subspecies. Advancements in laboratory techniques and continued investigations into rear-fanged venoms may uncover venoms that exhibit a great deal of complexity, and there is the distinct possibility of identifying novel venom compounds that exhibit unique pharmacological activities (Mackessy 2002; Saviola et al. 2014). Therefore, research examining rear-fanged venoms holds significant promise for novel compound discovery and is critical for refining our understanding of the evolutionary origin of venom systems in squamates.

## Venom evolution

Much debate centers on the evolutionary origin of venoms and the events that have promoted their toxic effects. It has been suggested that venoms evolved via the recruitment and duplication of genes with normal physiological function expressed elsewhere in the body (Fry 2005). This supposition has been supported in part by evidence of venom

gene homologs expressed in a variety of non-secretory tissues that encode toxins in the non-venomous Burmese python (Reyes-Velasco et al. 2015). Conversely, Hargreaves and colleagues (2014) have recently suggested that snake venoms evolved from pre-existing salivary proteins confined to the venom gland, and that venom toxins did not evolve from body proteins. At present, it appears most likely that ''venom protein'' homologs are broadly expressed throughout many tissues of both venomous and non-venomous snakes, but that the actual genes of proteins serving as venom toxins are overexpressed in the venom gland only (Reyes-Velasco et al. 2015; Junqueira-de-Azevedo et al. 2015). These data strongly indicate several points: (1) transcript-based sequences of oral gland transcripts/proteins only, derived from a variety of squamate reptiles (including oral gland secretions as venoms), should be interpreted with great caution, as these ''venom precursor'' genes are likely normal housekeeping genes; and (2) the ''raw material'' for the evolution of venom toxins are indeed derived from widely distributed genes with normal physiological roles (Fig. 2).

## F nction ers s biological role

The complex nature of venom is reflected in the presence of numerous venom compounds that not only have discrete functionalities, but also can act in concert with other venom components to promote a myriad of biological effects. Approximately 24 different protein families are represented in reptile venoms, with diverse pharmacologies, and approximately one-half of these are expressed in a single venom. It is interesting to note that some toxins within the same family may have widely disparate pharmacologies, as seen with $PLA_2$s that can exhibit neurotoxic, myotoxic, and anticoagulant effects. Many activities are represented (Table 1), and the end result of prey envenomation is a general and simultaneous dysregulation of numerous systems necessary for basic life functions. However, the biological role of an individual venom compound may significantly differ from its pharmacology, and a clear example of this can be seen with the disintegrin family of snake venom proteins. Disintegrins are small non-enzymatic proteins common in the venoms of viperid snakes (Calvete et al. 2005; Saviola et al. 2015a) that function by selectively blocking integrin receptors present in cell membranes (Eble et al. 2003; Bolás et al. 2014; Saviola et al. 2016). By blocking integrin $\alpha_{IIb}\beta_3$ on platelets, many disintegrins inhibit platelet aggregation and clot formation, complementing thrombin-like serine proteases and hemorrhagic SVMPs to promote the spread of other toxins throughout the envenomated prey. Interestingly, disintegrins also appear to be the ''relocator'' molecule in venom, allowing rattlesnakes to discriminate between envenomated and non-envenomated chemical cues through vomeronasal chemoreception (Saviola et al. 2013). Although the mechanism facilitating the changes in the chemical odor between envenomated and non-envenomated prey is still unknown, disintegrins clearly indicate how some venom proteins have unexpected, and often surprising, biological roles. In the following section, we discuss the function and biological roles of some common snake venom proteins, focusing on several that serve critical biological roles. Many of these compounds are well studied from the venoms of front-fanged snakes, and we will briefly address current work conducted on rear-fanged snake venoms and specific compounds that have been isolated from these venoms. A recent review presents a catalog of most known components of rear-fanged snake venoms (Junqueira-de-Azevedo et al. 2016).

### Snake venom metalloproteinases

It has long been known that many snake venoms exhibit significant proteolytic activity, and the observation that this activity can be abolished by metal chelators such as ethylenediaminetetraacetic acid (EDTA) or , -phenanthroline indicated that divalent cations are required for activity, giving rise to the name snake venom metalloproteinase (SVMP) (Satake et al., 1963). Zinc-dependent SVMPs are small to moderate sized enzymes that may consist of multiple domains, and along with ADAMs (a disintegrin and metalloproteinase), they comprise the M12 subfamily of metalloproteinases. SVMPs are thought to have evolved from early recruitment and modification of ADAM-like ancestors similar to ADAM 7, ADAM 28, and ADAMDEC1 (decysin) before the radiation of advanced snakes (Jia et al. 1996; Fry 2005; Casewell 2012). This hypothesis is largely supported by the presence of the largest SVMPs, the P-III class, in the venoms of species belonging to the families Viperidae, Elapidae, Atractaspididae, and Colubridae, although the expression levels of this enzyme can vary significantly

contains only this metalloproteinase domain. P-II SVMPs express an additional spacer region of amino acids carboxy to the metalloproteinase domain, followed by a disintegrin domain that is often proteolytically processed, liberating the disinte-

immobilization and tissue proteolysis, and they are hypothesized to assist in the pre-digestion of larger, heavy bodied prey items that are commonly consumed by many viperid and some colubrid snakes (Mackessy 1988, 2010). Conversely, elapids feed on more elongate ectothermic prey having higher surface-to-volume ratios, and their venoms typically show low to very low SVMP activity. However, there are conflicting reports on the contribution of venoms in prey digestion (see Thomas and Pough 1979; McCue 2007; Chu et al 2009); unfortunately, these studies used differing methodologies, so comparison across them is difficult.

SVMPs are broadly distributed in venoms of rear-fanged snakes (Hill and Mackessy 2000), and biochemical assays, as well as proteomic and transcriptomic approaches (Ching et al. 2006; Modahl et al. 2016a), demonstrate that SVMPs are one of the most abundant compounds in their venoms. P-III SVMPs were the most abundant and diverse toxin family present in the transcriptome of the Night Snake (*H [illegible]*) (McGivern et al. 2014), and Peichoto et al. (2012) identified SVMPs (based on activity and protein masses) in the venoms of *Ph[illegible]*, *P. ba[illegible]*, *P. [illegible]*, *H. [illegible]*, and *T[illegible]*. Individual SVMPs have been characterized from the venoms of *P. [illegible]* (Assakura et al. 1994), *Ph[illegible]* (Peichoto et al. 2007), and *A. [illegible]* (Weldon and Mackessy 2012), and all are P-III SVMPs. These isolated enzymes exhibited direct fibrin(ogen)olytic activity, cleaving either the A(α) or B(β)- chains of fibrinogen and/or fibrin; however, three of these SVMPs from the venom of *P. [illegible]* lacked the hemorrhagic activity (Assakura et al. 1994) that is common to many rear-fanged snake SVMPs.

## Phospholipase $A_2$

Phospholipase $A_2$ ($PLA_2$) enzymes constitute a major snake venom component and often contribute substantially to prey immobilization and capture. To date, $PLA_2$ activity, as well as isolated $PLA_2$s, have been identified and characterized from snakes belonging to the families Viperidae, Elapidae, and Colubridae (Kini 1997; Doley et al. 2010). Similar to SVMPs (and the majority of venom compounds), $PLA_2$ concentrations in venoms show tremendous phylogenetic variation. These enzymes share 40–99% amino acid sequence identity, as well as similarities in their three-dimensional structures, but despite their similarities, the pharmacological properties of individual $PLA_2$s may differ significantly. Phospholipases $A_2$ are among the most toxic and pharmacologically active venom compounds, and significant amounts of research have centered on these enzymes. Some elapids, such as the Central American Coral Snake (*M[illegible]*), produce venoms rich in $PLA_2$s, and approximately 48% of its venom proteome consists of this enzyme (Fernández et al. 2011). Among viperids, the Tiger Rattlesnake (*C[illegible]*) exhibits a simple yet extremely toxic venom, and approximately 66% of this venom consists of a homolog of the presynaptic β-neurotoxic $PLA_2$, crotoxin (Calvete et al. 2012). Similar in structure and function to crotoxin (discussed below), Mojave toxin is the primary toxic component in the venom of the Mojave Rattlesnake (*C. [illegible]*) and may comprise 45% of this species' venom in populations in southeastern Arizona, but it is completely absent in nearby north-central Arizona populations (Massey et al. 2012).

To date, little effort has been directed at rear-fanged snake venom $PLA_2$s; however, activity has been detected in venoms of several species (see Mackessy 2002; Huang and Mackessy 2004; Zelanis et al. 2010; Saviola et al. 2014). Trimorphin, a 13.9 kDa $PLA_2$ from the venom of the Sonoran Lyre Snake (*T[illegible]*), is the only $PLA_2$ purified and characterized from a rear-fanged snake venom. Analysis of the first 50 amino acid residues suggest that trimorphin is more closely related to $PLA_2$s from seas snakes and Australian elapids than to other terrestrial elapid or viperid $PLA_2$s (Huang and Mackessy 2004). Although the pharmacology of trimorphin has yet to be examined, this $PLA_2$ did exhibit significant and dose-dependent anti-parasitic (*Le[illegible]*) activity [illegible], with an $IC_{50}$ of 0.25 [illegible] M (Peichoto et al. 2011).

## Three-finger toxins

Three-finger toxins (3FTxs) were once thought to be unique to elapid venoms; however, the characterization of α-colubritoxin (Fry et al. 2003a), followed by venom protein isolation/characterization (Pawlak et al. 2006, 2009; Heyborne and Mackessy 2013) and transcriptomic and proteomic studies (Junqueira-de-Azevedo et al. 2006; Pahari et al 2007; McGivern et al. 2014; Modahl et al. 2016a), indicate that these toxins are more widely distributed among venoms of advanced snakes. 3FTxs have a highly conserved protein scaffold, with three finger-like loops stabilized by disulfide bonds protruding from their central core (Fig. 3). Differences in non-structural amino acid residues result in a multitude of pharmacological functions with a diversity of

Downloaded from http://icb.oxfordjournals.org/ at University of Colorado on October 24, 2016

biological roles, in spite of nearly identical crystal structure. Various 3FTxs have been shown to recognize an array of receptors, including nicotinic and muscarinic acetylcholine receptors (Kini and Doley 2010), L-type calcium channels, integrins (Kini 2002), coagulation factor VIIa (Banerjee et al. 2005), and β1/β2-adrenergic receptors (Rajagopalan et al. 2007). Based on these different ligand specificities, venom 3FTxs exhibit diverse biological activities in the form of neurotoxicity, cardiotoxicity, cytotoxicity, and anticoagulation effects (Hegde et al. 2010), some of which are discussed below.

### Cysteine-rich secretory proteins

Cysteine-rich secretory proteins (CRISPs) are a family of 20–30 kDa, non-enzymatic proteins widely distributed among reptile venoms. CRISPs display a high degree of conservation in their structure, due to high amino acid sequence similarities and 16 highly conserved cysteine residues that form eight disulfide bonds. Like the 3FTxs, despite their high degree of structural conservation, CRISPs show a variety of pharmacologies (reviewed by Heyborne and Mackessy 2010) including binding to cyclic nucleotide-gated ion channels (Brown et al. 1999; Yamazaki et al. 2002a), blocking vascular smooth muscle contraction (Yamazaki et al. 2002b), inhibiting $Ca^{2+}$ release from the sarcoplasmic reticulum (Morrissette et al. 1995), and blocking calcium currents in neurons (Nobile et al. 1996). CRISPs have been isolated from the venoms of Elapidae (Brown et al. 1999; Yamazaki et al. 2002a), Viperidae (Yamazaki et al. 2002b) and Colubridae (Peichoto et al. 2009), as well as from venom of the Mexican Beaded Lizard (*He. de, a h ,,.d h ,,.d* ; Mochca-Morales et al. 1990). A recent evaluation of CRISP molecular evolution suggests that they are under stronger positive selection in snakes than in lizards (Sunagar et al. 2012).

Among rear-fanged venomous snakes, CRISPs appear to be one of the more abundant venom proteins; however, characterization and functional data on these compounds are scarce. Patagonin, a 24.8 kDa CRISP isolated from the venom of *P. a a-g .e .*, exhibited unique necrotic activity toward murine gastrocnemius muscle at higher doses, although it did not induce edema or hemorrhage and it had no effect on the aggregation of human platelets or platelet-rich plasma (Peichoto et al. 2009). Patagonin did not exhibit proteolytic activity toward azocoll, azocasein, or fibrinogen. The widespread distribution of CRISPs among reptile venoms suggests a significant trophic role for these proteins, but their biological role(s) in envenomation are far from clear. It is possible that CRISPs may act in concert with other venom compounds to enhance venom lethality, but this possibility has not yet been explored.

**Fig. 3** Taxon-specific three-finger toxins from several species of rear-fanged snakes. Both toxins are potently lethal to lizards, but are non-toxic to mammals. (A) The Brown Treesnake (*Boiga irregularis*), an Old World colubrid, produces a venom containing approximately 10% irditoxin (**B**–backbone structure). (**C**) The Green Vinesnake (*Oxybelis fulgidus*) is a New World colubrid snake; fulgimotoxin (**D**) comprises approximately 35% of this species' venom. (**D**) Adapted from Heyborne and Mackessy (2013).

## Feeding strategies of enomo s snakes

Although venomous snakes utilize the same fundamental mechanism (envenomation) for limiting prey flight, differential strategies in prey envenomation and handling are seen both ontogenetically and between families. Viperids are generally considered ambush predators, and they use chemical cues in ambush site selection (Duvall et al. 1990; Clark 2004b) and visual-thermal cues to deliver what is most often a single envenomating strike (Hayes and Duvall 1991). Neonate rattlesnakes typically strike-and-hold small ectothermic prey (Mackessy 1988), whereas adults usually strike-and-release larger endothermic prey, returning to a reliance on chemical cues to relocate the envenomated prey that may wander from the attack site (Chiszar et al. 1977; Saviola et al. 2013). The release of prey following the envenomating strike allows snakes to avoid retaliation that may occur from larger, potentially dangerous prey items, yet requires relocating the meal using chemosensory searching once it has succumbed to venom (Chiszar et al. 1977; Parker and Kardong 2005; Saviola et al. 2013).

at University of Colorado on October 24, 2016

In contrast, the majority of snakes belonging to

*B. [illegible]* also demonstrates prey-specific lethality, with venoms being quite toxic toward chicken (*Ga[illegible]*; $LD_{50}$ = 1.75 μg/g) and lizard (*He[illegible]* geckos; $LD_{50}$ = 2.5 μg/g and *Ca[illegible]* skinks $LD_{50}$ = 4.5 μg/g) prey, whereas toxicity toward rodent prey (*M[illegible]*) was significantly lower ($LD_{50}$ = 31 μg/g; Mackessy et al. 2006). Irditoxin, a 17-kDa covalently linked heterodimeric 3FTx present in the venom of *B. [illegible]*, appears to be primarily responsible for toxic effects, and the purified toxin had $LD_{50}$ values of 0.22 μg/g and 0.55 μg/g for chicks and [illegible] respectively; [illegible]

canonical example of an elapid post-synaptic α-neurotoxin, binds very tightly to the nicotinic acetylcholine receptor α subunit of the vertebrate skeletal muscle motor endplate, resulting in rapid blockade of the ion channel and producing flaccid paralysis of

therefore appear to act as generalist toxins, potently lethal to divergent taxa of commonly utilized vertebrate prey.

### Venom metalloproteinases

Snake venom metalloproteinases (SVMPs) are found in the venoms of nearly all snakes, both front-fanged and rear-fanged, but some broad, global patterns of occurrence are observed. Vipers typically (but not always) have venoms rich in SVMPs, with 2–3 subtypes (PI-PIII) in a given venom, while elapids, and specifically marine species, typically (but not always) lack significant levels of these enzymes; they are also common to nearly all known rear-fanged snake venoms as well (Mackessy 2010). These enzymatic toxins catalyze the hydrolysis of many different proteins, particularly structural elements like collagen, elastin, and many proteins comprising connective tissues and the basal lamina/basement membranes of blood vessels and other epithelia (Fox and Serrano 2008; Oliveira et al. 2010). In this sense, their actions can be considered generalized, and structural degradation of prey tissues is a common manifestation of SVMP activity, in prey as diverse as crickets (Munekiyo and Mackessy 1998), lizards (Weldon and Mackessy 2012), and mammals (Moura-da-Silva et al 1996). However, a recent study of the specificity of PI, PII and PIII SVMPs toward basement membrane components, and proteomic analysis of wound exudates generated by individual SVMPs, indicated that although type IV collagen was a substrate common to all three subtypes, other structural elements were differentially affected (Herrera et al. 2015). Whether these specific actions apply to non-mammalian prey is unknown at present, but the three subtypes of SVMPs do appear to catalyze different types of structural damage.

## Summary and conclusions

### Why study rear-fanged snake venoms?

As venoms from more species of rear-fanged snakes are investigated, it is becoming apparent that many venom toxin families are shared among the advanced snakes, and some species show levels of venom complexity similar to those seen in venoms of elapids and viperids. However, relatively few rear-fanged snake species have been investigated, and sampling has often relied on specimen availability, rather than phylogenetically-directed sampling that is needed to encompass the diversity of venoms likely present

biochemistry of rear-fanged snakes reflects the complex evolutionary history of these species. Venoms have evolved from an apparently small selection of possible protein precursors (approximately 24 protein families out of thousands known: Wu et al. 2003; Bateman 2014), suggesting that venoms may have evolved only once in squamate reptiles. However, the disparate morphology of venom glands, the broad occurrence of venom ''homologs'' in non-oral gland tissues, different trends in venom composition among major clades and the long evolutionary history of squamates suggest that it is more parsimonious that venoms and venom delivery systems evolved several times in disparate squamate clades (Fig. 8).

It is important to keep in mind that we presently

## Acknowledgments

We thank Marymegen Daly and H. Lisle Gibbs for the invitation to participate in the 2016 SICB symposium Integrative and Comparative Biology of Venom, Steven D. Aird for his thoughtful and helpful editing/critique of the article, and two anonymous reviewers for their comments.

## Funding

Funding was provided by National Science Foundation (symposium: Daly/Gibbs) and a UNC Provost award (to S.P.M.).

## References

Aird SD, Kaiser II. 1985. Comparative studies on three rattlesnake toxins. Toxicon 23: 361–74.

Aird SD, Kaiser II, Lewis RV, Kruggel WG. 1985. Rattlesnake presynaptic neurotoxins: primary structure and evolution-

Sonoran Lyre Snake *T,* , *h d b c q a bda* (family Colubridae). Toxicon 44: 27–36.

Jia LG, Shimokawa KI, Bjarnason JB, Fox JW. 1996. Snake venom metalloproteinases: structure, function and relationship to the ADAMs family of proteins. Toxicon 34: 1269–76.

Junqueira-de-Azevedo ILM, Bastos CM, Ho PL, Luna MS, Yamanouye N, Casewell NR. 2015. Venom-related transcripts from *B h a a aca* tissues provide novel molecular insights into the production and evolution of snake venom. Mol Biol Evol 32: 754–66.

Junqueira-de-Azevedo ILM, Ching ATC, Carvalho E, Faria F, Nishiyama Jr MY, Ho PL, Diniz MRV. 2006. *Lache a* (Viperidae) cDNAs reveal diverging pit viper molecules and scaffolds typical of cobra (Elapidae) venoms: implications for snake toxin repertoire evolution. Genetics 173: 877–89.

Junqueira-de-Azevedo ILM, Campos PF, Ching ATC, Mackessy SP. 2016. Colubrid venom composition: an - omics perspective. Toxins 8: 230.

Kardong KV. 1982. Comparative study of changes in prey capture behavior of the cottonmouth (*Ag , d c*) and Egyptian cobra (*Na a ha e*). Copeia 1982: 337–43.

Kardong KV, Lavin-Murcio PA. 1993. Venom delivery of snakes as high–pressure and low–pressure systems. Copeia 1993: 644–50.

Moura-da-Silva AM, Theakston RD, Crampton JM. 1996. Evolution of disintegrin cysteine-rich and mammalian matrix-degrading metalloproteinases: gene duplication and divergence of a common ancestor rather than convergent evolution. J Mol Evol 43: 263–9.

Munekiyo SM, Mackessy SP. 1998. Effects of temperature and storage conditions on the electrophoretic, toxic and enzymatic stability of venom components. Comp Biochem Physiol B 119: 119–27.

Nirthanan S, Gwee MCE. 2004. Three-finger neurotoxins and the nicotinic acetylcholine receptor, forty years on. J Pharmacol Sci 94:1–17

Nobile M, Noceti F, Prestipino G, Possani LD. 1996. Helothermine, a lizard venom toxin, inhibits calcium current in cerebellar granules. Exp Brain Res 110:15–20.

Oliveira AK, Paes Leme AF, Asega AF, Camargo AC, Fox JW, Serrano SM. 2010. New insights into the structural elements involved in the skin haemorrhage induced by snake venom metalloproteinases. Thromb Haemost 104: 485–97.

Oliveira LD, Scartozzoni RR, Almeida-Santos SM, Jared C, Antoniazzi MM, Salomão MD. 2016. Morphology of