

Contents lists available at ScienceDirect

Toxicon

journal homepage: www.elsevier.com/locate/toxicon



Venom composition of adult Western Diamondback Rattlesnakes (Crotalus atrox) maintained under controlled diet and environmental conditions shows only minor changes

Christopher J. Rex[1], Stephen P. Mackessy

Department of Biological Sciences, 501 20th St., University of Northern Colorado, Greeley, CO, 80639-0017, USA

venom. In addition, understanding the extent of this variation has very practical applications, as e cacy of antivenoms could be compromised if high levels of antigenic novelty exist in di erent populations of the same species.

Ontogenetic and/or size-related changes in venom composition, in some cases paralleling a shift in the type/size of preferred prey, occurs in many species and is often most pronounced in viperids (Alape-Giron et al., 2008; Calvete et al., 2010; Daltry et al., 1996a; Gibbs et al., 2011; Mackessy, 1988, 1993; 2008; Mackessy et al., 2003, 2006; 2018; Zelanis et al., 2010). There has been speculation that these changes in gene expression levels have evolved in response to utilizing di erent prey types and sizes that snakes will encounter as they grow larger. Understanding whether venom composition is strictly genetically controlled or can be shaped by diet, environment or season will aid in comprehending how/why venom can sometimes be quite variable

150 × 4.6 mm, 5 µm; Grace Davison Discovery Sciences); total run time was 124 min (5% solvent B for 6 min, 5–15% solvent B over 3 min, 15–80% solvent B over 98 min, 80–100% solvent B over 2 min, 100% solvent B for 5 min, 100-5% solvent B for 2 min, and then 5% solvent B for 8 min). To conserve time and sample, two hundred µg of venom

presence/absence of venom proteins or peptides (in the range of 4000–84000 m/z) between first and last extractions, but again, minor differences between individual snakes were observed (Fig. 7).

3.6. Summary of qualitative analyses

Minimal differences between venom extractions within individual snakes and minor variation between individuals were observed (Table 1). There were no detectable differences in qualitative venom profiles between males and females or in relation to the time of the year

signi cantly (Table 1) between venom extractions one (311.1 ± 59.4), two (315.5 ± 50.9), three (312.3 ± 53.5), and four (323.6 ± 56.4; $p = 0.917$). There was no correlation between total length and LKLSP activity at the

47ble 5.7169 [(two)-419.1 a6.6 (at)-34CS0 cp.9

ls (ase/decrease between venom extractions one and four) in enzyme activity and protein concentration for all 16 P, oncentrations (Conc) show little change, LAAO and PDE show large positive increases in activity.

po s in enzyme activities 19 ver, no such changes in x, fact that LAAO, which iv yellow color, increased ol di erences in venom er vild-caught adult S. m. ith the greatest di er- suggesting

possibility that venom composition could also be in uenced by sex. Females produced venoms with ~8% higher AZO MPr activities than males, suggesting a sex-based dierence in this prominent venom protein family, but no other analyses revealed any signi cant di erences between sexes. Hypothetically, slightly lower AZO MPr activity could be related to heightened testosterone levels (and/or the resulting higher metabolic rates) that male C. atrox experience in comparison to females (Taylor and Denardo, 2005). Venom from all but one female showed a decrease in AZO MPr activity from rst to last ex- tatistically signi cant). Sex-

antivenom production or toxinological research will show signi cant di erences from their wild-caught counterparts. However, possible changes in venom composition that occur due to captivity and/or diet have been evaluated for only a small number of species, so it is possible that di erent taxa could show di erential e ects of captivity.

Conflicts of interest

The authors declare that there are no con icts of interest.

Acknowledgements

We thank Drs. T. Schountz and M. P. Thomas for excellent suggestions on C. J. Rex's Master's thesis, from which this manuscript was prepared. This research was supported by grants awarded by the University of Northern Colorado Graduate School and Graduate Student Association to CJR.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.toxicon.2019.03.027 .

Ethical statement

The authors hereby state that all procedures involving animals were conducted in a humane and ethical manner. All protocols were evaluated and approved (prior to initiating research) by the University of Northern Colorado Institutional Animal Care and Use Committee (UNC-IACUC).

References

Aird, S.D., 2008. Snake venom dipeptidyl peptidase IV: taxonomic distribution and quantitative variation. Comp. Biochem. Physiol. B Biochem. Mol. Biol. 150, 222 –228.

Alape-Giron, A., Sanz, L., Escolano, J., Flores-Diaz, M., Madrigal, M., Sasa, M., Calvete, J.J., 2008. Snake venomics of the lancehead pitviper Bothrops aspergeographic, individual, and ontogenetic variations. J. Proteome Res. 7, 3556–3571.

Beaupre, S.J., Duvall, D., 1998. Variation in oxygen consumption of the western diamondback rattlesnake (Crotalus atrox): implications for sexual size dimorphism. J.

.

Reid, H.A., Theakston, R.D.G., 1978. Changes in coagulation e